

Chiral quark models and their applications ^{*}

Jan O. Eeg[†]

*Department of Physics, University of Oslo,
P.O.Box 1048 Blindern, N-0316 Oslo, Norway*

Aksel Hiorth[‡]

RF-Rogaland Research, P.O.Box 8046, N-4068 Stavanger, Norway

We give an overview of chiral quark models, both for the pure light sector and the heavy-light sector. We describe how such models can be bosonized to obtain well known chiral Lagrangians which can be inferred from the symmetries of QCD alone. In addition, we can within these models calculate the coefficients of the various pieces of the chiral Lagrangians. We discuss a few applications of the models, in particular, $B - \bar{B}$ mixing and processes of the type $B \rightarrow D\bar{D}$, where D might be both pseudoscalar and vector. We suggest how the formalism might be extended to include light vectors (ρ, ω, K^*), and heavy to light transitions like $B \rightarrow \pi$.

I. INTRODUCTION

While the short distance (SD) effects in hadronic physics are well understood within perturbative quantum chromodynamics (pQCD), long distance (LD) effects have been hard to pin down. Even if quark models are not QCD itself, various QCD inspired quark models have been useful to make predictions for a limited class of problems. Lattice QCD and QCD sum rules are on more solid ground theoretically, but are in various cases not so easy to apply. In the light quark sector, low energy quantities have been studied in terms of the (extended) Nambu-Jona-Lasinio model (NJL)[1], and also the chiral quark model (χ QM)[2], which is the mean field approximation of NJL.

Within the χ QM, the light quarks (u, d, s) couple to the would be Goldstone octet mesons

^{*} Dedicated to the memory of Prof. Dubravko Tadić

[†]Electronic address: j.o.eeg@fys.uio.no

[‡]Electronic address: aksel.hiorth@rtf.no




FIG. 1: Energy hierarchy in interactions of elementary particles.

(K, π, η) in a chiral invariant way, such that all effects are in principle calculable in terms of physical quantities and a few model dependent parameters, namely the quark condensate, the gluon condensate, and the constituent quark mass [3, 4, 5]. More specific, one may calculate the coupling constants of chiral Lagrangians by integrating out the quarks by means of the χQM . In this way chiral quark models bridge between pQCD and chiral perturbation theory (χPT) as indicated in Fig. 1.

The ideas from the chiral quark model of the pure light sector [2, 3, 4, 5] has been extended to the sector involving a heavy quark (c or b) and thereby to heavy-light mesons [6]. Such models we name heavy-light chiral quark models ($\text{HL}\chi\text{QM}$). Also in this case, one may integrate out the light and heavy quarks and obtain chiral Lagrangians involving light and heavy mesons [7]. That is, we calculate the parameters of chiral Lagrangian terms, where the description of heavy mesons are in accordance with heavy quark effective field theory (HQEFT) [8]. In our approach [9] we extended the formalism of [6] to include gluon (vacuum) condensates.

One important motivation for the inclusion of gluon condensates is the possibility to estimate non-factorizable (colour suppressed) contributions in non-leptonic decays. For instance, $K - \bar{K}$ -mixing and the $\Delta I = 1/2$ rule for $K \rightarrow 2\pi$ can be understood in a reasonable way within the χ QM [4, 5] including gluon condensates. Especially, the suppression of the $I = 2$ amplitude found for $K \rightarrow 2\pi$ is also in agreement with generalized factorization [10]. Furthermore, it allows us for instance to consider decays where the gluonic aspect of η' is relevant [11], and some aspects of D -meson decays [12]. The most important application is to calculate non-factorizable contributions to $B - \bar{B}$ -mixing [13], where our approach includes $1/m_b$ corrections and chiral corrections both from loops and counterterms. Also processes of the type $B \rightarrow D\bar{D}$ are calculable [14]. It should be emphasized that the HL χ QM can not, -in its present form, be used for heavy to light transitions like $B \rightarrow \pi K$, where QCD factorization [15] or soft collinear theory(SCET) [16] is often applied. Still, in the last section, we suggest how an extension to this case might be performed. We also suggest how the χ QM might be extended to include light vectors (ρ, ω, K^*).

II. CHIRAL PERTURBATION THEORY

A. The pure light sector

Quarks are the fundamental hadronic matter. However, the particles we observe are those built out of them: baryons and mesons. In the sector of the lowest mass pseudoscalar mesons (the would-be Goldstone bosons: π , K and η) the interactions can be described in terms of an effective theory, the chiral Lagrangian, that includes only these states. The chiral Lagrangian and chiral perturbation theory (χ PT) [17, 18] provide a faithful representation of this sector of the Standard Model after the quark and gluon degrees of freedom have been integrated out. The form of this effective field theory and all its possible terms are determined by $SU_L(3) \times SU_R(3)$ chiral invariance and Lorentz invariance. Terms which explicitly break chiral invariance are introduced in terms of the quark mass matrix $\mathcal{M}_q$.

The strong chiral lagrangian is completely fixed to the leading order in momenta by symmetry requirements and the Goldstone boson's decay amplitudes:

$$\mathcal{L}_{\text{strong}}^{(2)} = \frac{f^2}{4} \text{Tr} (D_\mu \Sigma D^\mu \Sigma^\dagger) + \frac{f^2}{2} B_0 \text{Tr} (\mathcal{M}_q \Sigma^\dagger + \Sigma \mathcal{M}_q^\dagger), \quad (1)$$

where the covariant derivative D^μ contains the photon field, and $\mathcal{M}_q = \text{diag}[m_u, m_d, m_s]$.

The quantity B_0 is defined by $\langle \bar{q}_i q_j \rangle = -f^2 B_0 \delta_{ij}$, where

$$(m_s + m_d) \langle \bar{q} q \rangle = -f_K^2 m_K^2, \quad (m_u + m_d) \langle \bar{q} q \rangle = -f_\pi^2 m_\pi^2, \quad (2)$$

in the PCAC limit. The quantity $\langle \bar{q} q \rangle$ is the quark condensate, being of order $(-240 \text{ MeV})^3$.

The $SU_L(3) \times SU_R(3)$ field Σ contains the pseudoscalar octet Π :

$$\Sigma \equiv \exp \left(\frac{2i}{f} \Pi \right), \quad \Pi = \frac{\lambda^a}{2} \phi^a(x) = \frac{1}{\sqrt{2}} \begin{bmatrix} \frac{\pi^0}{\sqrt{2}} + \frac{\eta_8}{\sqrt{6}} & \pi^+ & K^+ \\ \pi^- & -\frac{\pi^0}{\sqrt{2}} + \frac{\eta_8}{\sqrt{6}} & K^0 \\ K^- & \bar{K}^0 & -\frac{2}{\sqrt{6}} \eta_8 \end{bmatrix}. \quad (3)$$

The quantity f is, to lowest order, identified with the pion decay constant f_π (and equal to f_K before chiral loops are introduced).

When the matrix Σ is expanded in powers of f^{-1} , the zeroth order term obtained from (1) is the free Klein-Gordon Lagrangian for the pseudoscalar particles. From this Lagrangian one might deduce the (left-handed) current

$$J_\mu^n = -i \frac{f^2}{2} \text{Tr} [\lambda^n \Sigma D^\mu \Sigma^\dagger], \quad (4)$$

where n is a flavour octet index and λ_n a $SU(3)$ flavour matrix.

For the next-to-leading order Lagrangian $\mathcal{L}_{\text{strong}}^{(4)}$ there are ten terms and thereby ten coefficients L_i to be determined [18] either experimentally or by means of some model. Some of these play an important role in the physics of ϵ' in $K \rightarrow 2\pi$ decays[19]. As examples, we display the L_5 and L_8 terms in governing much of the penguin physics:

$$L_5 B_0 \text{Tr} [D_\mu \Sigma^\dagger D^\mu \Sigma (\mathcal{M}_q^\dagger \Sigma + \Sigma^\dagger \mathcal{M}_q)] , \quad (5)$$

and

$$L_8 B_0 \text{Tr} [\mathcal{M}_q^\dagger \Sigma \mathcal{M}_q^\dagger \Sigma + \mathcal{M}_q \Sigma^\dagger \mathcal{M}_q \Sigma^\dagger] . \quad (6)$$

Under the action of the elements V_R and V_L of the chiral group $SU_R(3) \times SU_L(3)$, the field Σ transforms as:

$$\Sigma \rightarrow V_L \Sigma V_R^\dagger, \quad (7)$$

and accordingly for the conjugated fields. Formally, $\mathcal{M}_q$ is given the same transformation properties as Σ , and $\mathcal{M}_q^\dagger$ as $\Sigma^\dagger$.

B. The heavy light sector

The strong chiral Lagrangian for the heavy light sector is [7, 20]:

$$\begin{aligned} \mathcal{L}_{Str} &= \mp Tr \left[\overline{H_{vk}^{(\pm)}} (i v \cdot \mathcal{D}_{hk}) H_{vh}^{(\pm)} \right] - g_A Tr \left[\overline{H_{vk}^{(\pm)}} H_{vh}^{(\pm)} \gamma_\mu \gamma_5 \mathcal{A}_{hk}^\mu \right] \\ &+ 2\lambda_1 Tr \left[\overline{H_{vk}^{(\pm)}} H_{vh}^{(\pm)} (\widetilde{M}_q^V)_{hk} \right] + \frac{e\beta}{4} Tr \left[\overline{H_{vk}^{(\pm)}} H_{vh}^{(\pm)} \sigma \cdot F(Q_q^\xi)_{hk} \right] + \dots \end{aligned} \quad (8)$$

where k, h are $SU(3)$ triplet indices, and v is the velocity of the heavy meson. The ellipses indicate other terms (of higher order, say), and $i\mathcal{D}_{hk}^\mu = \delta_{hk} D^\mu + \mathcal{V}_{hk}^\mu$. Moreover, $Q_q^\xi = (\xi^\dagger Q_q \xi + \xi Q_q \xi^\dagger)/2$, where Q_q is the $SU(3)$ charge matrix for light quarks, $Q_q = \text{diag}(2/3, -1/3, -1/3)$, and F is the electromagnetic field tensor. $H_{vk}^{(\pm)}$ is the heavy meson field containing a spin zero and spin one boson:

$$\begin{aligned} H_{vk}^{(\pm)} &\equiv P_\pm(v) (P_k^{(\pm)\mu} \gamma_\mu - i P_k^{(\pm)5} \gamma_5) , \\ \overline{H_{vk}^{(\pm)}} &= \gamma^0 (H_{vk}^{(\pm)})^\dagger \gamma^0 = \left[(P_k^{(\pm)\mu})^\dagger \gamma_\mu - i (P_k^{(\pm)5})^\dagger \gamma_5 \right] P_\pm , \end{aligned} \quad (9)$$

where

$$P_\pm(v) = (1 \pm \gamma \cdot v)/2 \quad (10)$$

are projection operators. The fields $P^{(\pm)5}(P^{(\pm)\mu})$ represent heavy-light mesons, $0^-(1^-)$, with velocity v . The signs $\pm$ refers to particles and anti-particles respectively, and will sometimes be omitted in the following when unnecessary.

The vector and axial vector fields $\mathcal{V}_\mu$ and $\mathcal{A}_\mu$ are given by:

$$\mathcal{V}_\mu \equiv \frac{i}{2} (\xi^\dagger \partial_\mu \xi + \xi \partial_\mu \xi^\dagger) \quad ; \quad \mathcal{A}_\mu \equiv -\frac{i}{2} (\xi^\dagger \partial_\mu \xi - \xi \partial_\mu \xi^\dagger) . \quad (11)$$

The fields ξ and H_v transform as

$$\xi \rightarrow U \xi V_R^\dagger = V_L \xi U^\dagger \quad , \quad H_v \rightarrow H_v U^\dagger , \quad (12)$$

where $U \in SU(3)_V$, the unbroken symmetry.

The vector and axial fields transform as

$$\mathcal{V}_\mu \rightarrow U \mathcal{V}_\mu U^\dagger + i U \partial_\mu U^\dagger \quad , \quad \mathcal{A}_\mu \rightarrow U \mathcal{A}_\mu U^\dagger . \quad (13)$$

The vector field $\mathcal{V}^\mu$ is seen to transform as a gauge field under local $SU(3)_V$, and can only appear in combination with a derivative as a covariant derivative ($i\partial^\mu + \mathcal{V}^\mu$). The quantity $\widetilde{M}_q^V$ (as well as the orthogonal combination $\widetilde{M}_q^A$) is related to the current mass term:

$$\widetilde{M}_q^V \equiv \frac{1}{2} (\xi^\dagger \mathcal{M}_q \xi^\dagger + \xi \mathcal{M}_q^\dagger \xi) \quad ; \quad \widetilde{M}_q^A \equiv -\frac{1}{2} (\xi^\dagger \mathcal{M}_q \xi^\dagger - \xi \mathcal{M}_q^\dagger \xi) . \quad (14)$$

The heavy-light weak current, to zeroth order in $1/m_Q$ and chiral counting, is represented by:

$$J_k^\alpha(0) = \frac{\alpha_H}{2} Tr \left[\xi_{hk}^\dagger \Gamma^\alpha H_{vh} \right] , \quad (15)$$

and under $SU(3)_L$ it transforms as

$$J_k^\alpha \rightarrow J_h^\alpha \left(V_L^\dagger \right)_{hk} . \quad (16)$$

This current has also (counter) terms, of higher order in the chiral counting, needed to make the chiral loops finite:

$$J_k^\mu(\mathcal{M}) = \frac{\omega_1}{2} Tr[\xi_{hk}^\dagger \Gamma^\mu H_{vl} \widetilde{\mathcal{M}}_{lh}^V] + \frac{\omega'_1}{2} Tr[\xi_{kh}^\dagger \Gamma^\mu H_{vh}] \widetilde{\mathcal{M}}_{ll}^V , \quad (17)$$

where the parameters ω_1 and ω'_1 are commented on in section IV-C. To leading order, $\Gamma^\alpha = \gamma^\alpha L$, where L is the left - handed projector in Dirac space, $L = (1 - \gamma_5)/2$. However, this is slightly modified by perturbative QCD for μ below m_Q , which gives [8]

$$\Gamma^\alpha \equiv C_\gamma(\mu) \gamma^\alpha L + C_v(\mu) v^\alpha R , \quad (18)$$

where R is the right - handed projector, $R = (1 + \gamma_5)/2$. The coefficients $C_{\gamma,v}(\mu)$ are determined by QCD renormalization for $\mu < m_Q$. They have been calculated to NLO and the result is the same in MS and $\overline{MS}$ scheme[21]. (C_γ is close to one and C_v is rather small). Corrections to the weak current of order $1/m_Q$ will be discussed in section V.

Before closing this section, we write down the bosonized $b \rightarrow c$ transition current in terms of the heavy fields

$$\overline{Q_{v_b}^{(+)}} \gamma^\mu L Q_{v_c}^{(+)} \longrightarrow -\zeta(\omega) Tr \left[\overline{H_c^{(+)}} \gamma^\alpha L H_b^{(+)} \right] , \quad (19)$$

where $\zeta(\omega)$ is the Isgur-Wise function for the $\bar{B} \rightarrow D$ transition [22]. The indices on the heavy fields here refer to the b - and c -quarks with velocities v_b and v_c , with $\omega \equiv v_b \cdot v_c$. The current for $D\bar{D}$ production is :

$$\overline{Q_{v_c}^{(+)}} \gamma^\mu L Q_{\bar{v}}^{(-)} \longrightarrow -\zeta(-\lambda) Tr \left[\overline{H_c^{(+)}} \gamma^\alpha L H_{\bar{c}}^{(-)} \right] , \quad (20)$$

where the Isgur-Wise function $\zeta(-\lambda)$ is (in general) complex. We have $\lambda = v_c \cdot \bar{v}$, where $\bar{v}$ is the velocity of $\bar{c}$.

III. THE CHIRAL QUARK MODEL (χ QM)

A. The Lagrangians for χ QM

The light quark sector is described by the chiral quark model (χ QM), having a standard QCD term and a term describing interactions between quarks and (Goldstone) mesons [1, 2, 3, 4, 5]:

$$\mathcal{L}_{\chi QM} = \bar{q}_L i\gamma \cdot D q_L + \bar{q}_R i\gamma \cdot D q_R - \bar{q}_L \mathcal{M}_q q_R - \bar{q}_R \mathcal{M}_q^\dagger q_L - m(\bar{q}_R \Sigma^\dagger q_L + \bar{q}_L \Sigma q_R) , \quad (21)$$

where m is the ($SU(3)$ - invariant) constituent quark mass for light quarks $q^T = (u, d, s)$. The left- and right-handed projections q_L and q_R are transforming after $SU(3)_L$ and $SU(3)_R$ respectively:

$$q_L \rightarrow V_L q_L \quad \text{and} \quad q_R \rightarrow V_R q_R . \quad (22)$$

From (21) we deduce the Feynman rules. For instance, the $Pq\bar{q}$ coupling is ($m\gamma_5/f$) times some $SU(3)$ factor (P is a pseudoscalar meson π, K, η). From such Feynman rules, and including the quark propagator $S(p) = (\gamma \cdot p - M_q)^{-1}$, we can calculate amplitudes for, say, $\pi - \pi$ scattering in the strong sector. Here $M_q = m + m_q$ is the total mass. Alternatively one might keep only the constituent mass m in the propagator, and take the current mass m_q as a coupling. Including also the Feynman rules for weak vertices, one might calculate [3, 5] amplitudes for non-leptonic decays in terms of quark loops representing f_π and the semileptonic form factors $f_\pm$, but also for more complicated cases.

Also, as a more exotic example, one may calculate the effect of the electroweak $s \rightarrow d$ self-energy transition contribution to $K \rightarrow 2\pi$ as shown in Fig. 2. This is an off-shell effect which vanishes in the free quark case, but is non-zero for bound quarks and proportional to m within our framework [3].

Chiral Lagrangians, either in the pure strong sector or for non-leptonic decays, are however easier to obtain in a more transparent way within the “rotated version” of the χ QM with flavour rotated quark fields χ given by:

$$\chi_L = \xi^\dagger q_L \quad ; \quad \chi_R = \xi q_R \quad ; \quad \xi \cdot \xi = \Sigma . \quad (23)$$

The constituent quark fields χ_L and χ_R transform in a simple way under $SU(3)_V$:

$$\chi_L \rightarrow U \chi_L \quad , \quad \chi_R \rightarrow U \chi_R . \quad (24)$$




FIG. 2: Contribution to the process $K \rightarrow 2\pi$ from the non-diagonal $s \rightarrow d$ transition.

In the rotated version, the chiral interactions are rotated into the kinetic term while the interaction term proportional to m in (21) become a pure (constituent) mass term [2, 5]:

$$\mathcal{L}_{\chi QM} = \bar{\chi} [\gamma^\mu (iD_\mu + \mathcal{V}_\mu + \gamma_5 \mathcal{A}_\mu) - m] \chi - \bar{\chi} \widetilde{M}_q \chi, \quad (25)$$

which is manifestly invariant under $SU(3)_V$. Moreover,

$$\widetilde{M}_q \equiv \widetilde{M}_q^V + \widetilde{M}_q^A \gamma_5, \quad (26)$$

where $\widetilde{M}_q^{V,A}$ are given in (14).

In the light sector, the various pieces of the strong Lagrangian in section II-A can now be obtained by integrating out the constituent quark fields χ , and these pieces can be written in terms of the fields $\mathcal{A}_\mu$, $\widetilde{M}_q^V$ and $\widetilde{M}_q^A$. This can easily be seen by using the relation

$$\mathcal{A}_\mu = -\frac{1}{2i} \xi (D_\mu \Sigma^\dagger) \xi = +\frac{1}{2i} \xi^\dagger (D_\mu \Sigma) \xi^\dagger. \quad (27)$$

In our model, the hard gluons are integrated out and we are left with soft gluonic degrees of freedom. These gluons can be described using the external field technique, and their effect will be parameterized by vacuum expectation values, i.e. the gluon condensate $\langle \frac{\alpha_s}{\pi} G^2 \rangle$. Gluon condensates with higher dimension could also be included, but we truncate the expansion by keeping only the condensate with lowest dimension.

When calculating the soft gluon effects in terms of the gluon condensate, we follow the prescription given in [23]. The calculation is easily carried out in the Fock - Schwinger gauge. In this gauge one can expand the gluon field as :

$$A_\mu^a(k) = -\frac{i(2\pi)^4}{2} G_{\rho\mu}^a(0) \frac{\partial}{\partial k_\rho} \delta^{(4)}(k) + \dots. \quad (28)$$

$$- \frac{1}{2} g_s t^a \gamma^\mu G_{\mu\rho}(0) \left. \frac{\partial}{\partial k^\rho} \right|_{k^\rho=0} + \dots$$

FIG. 3: Feynmanrule for the light quark -soft gluon vertex.

In some simple cases one may also use the light quark propagator in a gluonic background (to first order in the gluon field):

$$S_1(p, G) = -\frac{g_s}{4} G_{\alpha\beta}^b t^b [\sigma^{\alpha\beta} (\gamma \cdot p + m) + (\gamma \cdot p + m) \sigma^{\alpha\beta}] (p^2 - m^2)^{-2}, \quad (29)$$

where g_s is the strong coupling constant, a and b are colour octet indices, and t^a are the colour matrices. In general one should stick to the prescription in [23] in order to get correct results. Since each vertex in a Feynman diagram is accomplished with an integration we get the Feynman rule given in Fig. 3. The gluon condensate contributions are obtained by the replacement

$$g_s^2 G_{\mu\nu}^a G_{\alpha\beta}^b \rightarrow \frac{4\pi^2}{(N_c^2 - 1)} \delta^{ab} \langle \frac{\alpha_s}{\pi} G^2 \rangle \frac{1}{12} (g_{\mu\alpha} g_{\nu\beta} - g_{\mu\beta} g_{\nu\alpha}). \quad (30)$$

.

B. Bosonization of the χ QM

The Lagrangians (21) or (25) from the previous section can now be used for bosonization, i.e. to integrate out the quark fields. This can be done in the path integral formalism, or as we do here, by expanding in terms of Feynman diagrams. Within the χ QM, with Feynman rules obtained from (21) one may calculate the simple quark loop amplitude for $\pi \rightarrow W$ which defines f (the bare f_π) in terms of a logarithmically divergent integral I_2 times the coupling $\sim m/f$. Including also the gluon condensate contribution one obtains [3, 4, 5]:

$$f^2 = -i4m^2 N_c I_2 + \frac{1}{24m^2} \langle \frac{\alpha_s}{\pi} G^2 \rangle, \quad (31)$$

where I_2 is the following logarithmic divergent integral (d is the dimension of space within dimensional regularization):

$$I_2 \equiv \int \frac{d^d k}{(2\pi)^d} \frac{1}{(k^2 - m^2)^2}. \quad (32)$$

Equivalently, one may obtain the (kinetic part of the) strong Lagrangian in (1) by attaching two axial fields $\mathcal{A}_\mu$ to a vacuum polarization like quark loop diagram by using (25). Then one obtains:

$$i\mathcal{L}_{str}^{(2)} = -N_c \int \frac{d^d p}{(2\pi)^d} Tr [(\gamma_\sigma \gamma_5 \mathcal{A}^\sigma) S(p) (\gamma_\rho \gamma_5 \mathcal{A}^\rho) S(p)] = f^2 Tr [\mathcal{A}_\mu \mathcal{A}^\mu] , \quad (33)$$

where the trace is both in flavor and Dirac spaces (a similar diagram with gluons should also be added). This is easily seen by using the relation (27), provided f^2 is given by (31). The eq. (33) give the Lagrangian (1) in the light sector by applying (27).

The quark condensate is:

$$\langle \bar{q}q \rangle = -iN_c Tr \int \frac{d^d p}{(2\pi)^d} S(p) = -4imN_c I_1 - \frac{1}{12m} \langle \frac{\alpha_s}{\pi} G^2 \rangle , \quad (34)$$

where I_1 is the quadratically divergent integral

$$I_1 \equiv \int \frac{d^d k}{(2\pi)^d} \frac{1}{k^2 - m^2} . \quad (35)$$

Here the propagator S has to be understood at the one in the gluon field up to second order and the a priori divergent integrals $I_{1,2}$ have to be interpreted as the regularized ones. The physical values of $I_{1,2}$ are determined by the physical values of f and $\langle \bar{q}q \rangle$. In general, by coupling the fields $\mathcal{A}_\mu$ and $\widetilde{M}_q^{V,A}$ to quark loops, the chiral Lagrangian terms and their coefficients within the light sector can be obtained.

Similarly, we may bosonize the weak currents. The left handed current can be written

$$\bar{q}_L \gamma^\mu \lambda^n q_L = \bar{\chi}_L \gamma^\mu \Lambda^n \chi_L ; \quad \Lambda^n \equiv \xi^\dagger \lambda^n \xi . \quad (36)$$

The lowest order term $\mathcal{O}(p)$ is obtained when the vertex Λ^n from (36) and axial vertex ($\sim \mathcal{A}_\mu$) from (25) are entering a quark loop (see Fig. 4):

$$j_\mu^n(\mathcal{A}) = -iN_c \int \frac{d^d p}{(2\pi)^d} Tr [(\gamma_\mu L \Lambda^n) S(p) (\gamma_\sigma \gamma_5 \mathcal{A}^\sigma) S(p)] \sim Tr [\Lambda^n \mathcal{A}_\mu] , \quad (37)$$

which coincides with (4) when (27) is used.

As a more non-trivial example, to obtain a non-zero non-factorizable contribution to $D^0 \rightarrow K^0 \bar{K}^0$ at tree level, one has to consider the coloured current $j_\mu^{n,a}$ to $\mathcal{O}(p^3)$, involving insertions of the “mass fields” $\widetilde{M}_q$ in (26) [12]. (This coloured current is obtained by Fierz transformations of the relevant four quark operator). From Fig. 5, one obtains the contribution:




FIG. 4: Feynman diagram for bosonization of the left-handed current to order $\mathcal{O}(p)$.




FIG. 5: Diagram for bosonization of the colour current to $\mathcal{O}(p^3)$.

$$j_\mu^{n,a}(\text{Fig. 5}) = i \int \frac{d^d p}{(2\pi)^d} \text{Tr} \left[(\gamma_\mu L \Lambda^n t^a) S(p) (\gamma_\sigma \gamma_5 \mathcal{A}^\sigma) S(p) \widetilde{M}_q S_1(p, G) \right], \quad (38)$$

Summing all six diagrams with permutated vertices compared to the one in Fig. 5 we obtain in total :

$$j_\mu^{n,a}(G, \xi, \mathcal{A}, \widetilde{\mathcal{M}}_q) = \frac{g_s}{12m} \frac{1}{16\pi^2} G^{a,\kappa\lambda} \left[i\varepsilon_{\mu\rho\kappa\lambda} T_\varepsilon^{n,\rho} + (g_{\mu\kappa} g_{\rho\lambda} - g_{\mu\lambda} g_{\rho\kappa}) T_g^{n,\rho} \right], \quad (39)$$

where (we have used the analytical computer program FORM [24])

$$T_\varepsilon^{n,\rho} = 4 S_\rho^K - 3(S_\rho^L + S_\rho^R), \quad T_g^{n,\rho} = S_\rho^L - S_\rho^R. \quad (40)$$

The S' 's are chiral Lagrangian terms:

$$\begin{aligned} S_\rho^L &\equiv \text{Tr} \left[\Lambda^n \mathcal{A}_\rho \widetilde{M}_q^L \right] = \frac{1}{2i} \text{Tr} \left[\lambda^n (D_\rho \Sigma) \mathcal{M}_q^\dagger \right], \\ S_\rho^R &\equiv \text{Tr} \left[\Lambda^n \widetilde{M}_q^R \mathcal{A}^\rho \right] = \frac{-1}{2i} \text{Tr} \left[\lambda^n \mathcal{M}_q (D_\rho \Sigma^\dagger) \right], \\ S_\rho^K &\equiv \frac{1}{2} \text{Tr} \left[\Lambda^n \left(\mathcal{A}^\rho \widetilde{M}_q^R + \widetilde{M}_q^L \mathcal{A}^\rho \right) \right] \\ &= \frac{1}{4i} \text{Tr} \left[\lambda^n \left((D_\rho \Sigma) \Sigma^\dagger \mathcal{M}_q \Sigma^\dagger - \Sigma \mathcal{M}_q^\dagger \Sigma (D_\rho \Sigma^\dagger) \right) \right]. \end{aligned} \quad (41)$$

The current (39) has to be combined with the left-handed colour current for D -meson decay later given in (90) to obtain a contribution to $D^0 \rightarrow K^0 \bar{K}^0$ [12].

IV. THE HEAVY - LIGHT CHIRAL QUARK MODEL (HL χ QM)

A. The Lagrangian for HL χ QM

Our starting point is the following Lagrangian containing both quark and meson fields:

$$\mathcal{L} = \mathcal{L}_{HQEFT} + \mathcal{L}_{\chi QM} + \mathcal{L}_{Int} , \quad (42)$$

where [8]

$$\mathcal{L}_{HQEFT} = \pm \overline{Q_v^{(\pm)}} i v \cdot D Q_v^{(\pm)} + \frac{1}{2m_Q} \overline{Q_v^{(\pm)}} \left(-C_M \frac{g_s}{2} \sigma \cdot G + (iD_\perp)_{\text{eff}}^2 \right) Q_v^{(\pm)} + \mathcal{O}(m_Q^{-2}) \quad (43)$$

is the Lagrangian for heavy quark effective field theory (HQEFT). The heavy quark field $Q_v^{(+)}$ annihilates a heavy quark with velocity v and mass m_Q . Similarly, $Q_v^{(-)}$ annihilates a heavy anti-quark. Moreover, D_μ is the covariant derivative containing the gluon field (eventually also the photon field), and $\sigma \cdot G = \sigma^{\mu\nu} G_{\mu\nu}^a t^a$, where $\sigma^{\mu\nu} = i[\gamma^\mu, \gamma^\nu]/2$, and $G_{\mu\nu}^a$ is the gluonic field tensor. This chromo-magnetic term has a factor C_M , being one at tree level, but slightly modified by perturbative QCD. (When the covariant derivative also contains the photon field, there is also a corresponding magnetic term $\sim \sigma \cdot F$, where $F^{\mu\nu}$ is the electromagnetic tensor). Furthermore, $(iD_\perp)_{\text{eff}}^2 = C_D (iD)^2 - C_K (iv \cdot D)^2$. At tree level, $C_D = C_K = 1$. Here, C_D is not modified by perturbative QCD, while C_K is different from one due to perturbative QCD corrections for $\mu < m_Q$ [25]. We observe that soft gluons coupling to a heavy quark is suppressed by $1/m_Q$, since to leading order the vertex is proportional to $v_\mu v_\nu G^{a\mu\nu} = 0$, v_μ being the heavy quark velocity.

In the heavy - light case, the generalization of the meson - quark interactions in the pure light sector χ QM is given by the following $SU(3)$ invariant Lagrangian:

$$\mathcal{L}_{Int} = -G_H \left[\overline{\chi_k} \overline{H_{vk}^{(\pm)}} Q_v^{(\pm)} + \overline{Q_v^{(\pm)}} H_{vk}^{(\pm)} \chi_k \right] , \quad (44)$$

where k is a triplet $SU(3)$ - index and G_H is a coupling constant. Note that in [6], $G_H = 1$ is used. However, in that case one used a renormalization factor for the heavy meson fields H_v , which is equivalent. The interaction Lagrangian (44) can, as for the χ QM, be obtained from a NJL model. This has been done in [6] (-as for the light sector [1]).

B. Bosonization within the HL χ QM

The interaction term $\mathcal{L}_{Int}$ in (44) can now be used to bosonize the model, i.e. integrate out the quark fields. This can be done in terms of Feynman diagrams as we do here, by attaching the external fields $H_v^a, \overline{H}_v^a, \mathcal{V}^\mu, \mathcal{A}^\mu$ and $\widetilde{M}_q^{V,A}$ of section II-B to quark loops, and using (25) and (44). In this way one obtains the strong chiral Lagrangian (8) and terms of higher order in the heavy light sector. Some of the loop integrals will be divergent and have to be related to physical parameters, as for the pure light sector [2, 3, 4, 5]. As the pure light sector is a part of our model, we have to keep the relations in (31) and (34) from the pure light sector within the heavy light case studied here. The $1/m_Q$ terms will not be discussed in this section, but will be considered later in section V.

From the diagrams in Fig. 6 we obtain the identification for the kinetic term, which by Ward identities is the same as for the term with the vector field $\mathcal{V}_\mu$ attached to the light quark:

$$-iG_H^2 N_c (I_{3/2} + 2mI_2 + i \frac{(8-3\pi)}{384N_c m^3} \langle \frac{\alpha_s}{\pi} G^2 \rangle) = 1, \quad (45)$$

where I_2 is given in (32) and

$$I_{3/2} \equiv \int \frac{d^d k}{(2\pi)^d} \frac{1}{(v \cdot k)(k^2 - m^2)}, \quad (46)$$

which formally depends on v^2 which is equal to one. From the same diagram, with the axial field $\mathcal{A}_\mu$ attached, we obtain the following identification for the axial vector coupling g_A :

$$g_A \equiv iG_H^2 N_c \left(\frac{1}{3} I_{3/2} - 2mI_2 - i \frac{m}{12\pi} - i \frac{(8-3\pi)}{384N_c m^3} \langle \frac{\alpha_s}{\pi} G^2 \rangle \right), \quad (47)$$

As I_1 and I_2 are related to the quark condensate and f_π respectively, the (formally) linearly divergent integral $I_{3/2}$ is related to $\delta g_A \equiv 1 - g_A$, which is found by eliminating I_2 from eqs. (45) and (47):

$$\delta g_A = -\frac{4}{3} iG_H^2 N_c \left(I_{3/2} - i \frac{m}{16\pi} \right). \quad (48)$$

Note that the gluon condensate drops out here. Within a primitive cut-off regularization, $I_{3/2}$ is (in the leading approximation) proportional to the cut-off in first power:

$$I_{3/2} = i \frac{\Lambda}{16\pi} \left(1 + \mathcal{O}\left(\frac{m}{\Lambda}\right) \right), \quad (49)$$

where the cut-off Λ , is of the same order as the chiral symmetry breaking scale Λ_χ . In contrast, $I_{3/2}$ is finite and proportional to m in dimensional regularization. Note that the




FIG. 6: Bosonization in the strong sector to obtain eq. (8).

cut-off Λ is only used in qualitative considerations here and in subsection IV-D. Anyway, $I_{3/2}$ is determined by the physical value of g_A .

When attaching $\widetilde{M}_q^V$ like in Fig. 6 instead of vector or axial vector fields one finds for the mass correction term in (8):

$$2\lambda_1 \equiv iG_H^2 N_c (I_{3/2} - 2mI_2 - i\frac{m}{8\pi} - \frac{i(3\pi - 4)}{192N_c m^3} \langle \frac{\alpha_s}{\pi} G^2 \rangle) . \quad (50)$$

The electromagnetic β term in (8) is obtained by considering diagrams like those in figure 6, but with the vector and axial vector fields $\mathcal{V}_\mu$ or $\mathcal{A}_\mu$ replaced by a photon field tensor:

$$\beta = \frac{G_H^2}{2} \left\{ -4iN_c I_2 + \frac{N_c}{4\pi} - \left(\frac{32 + 3\pi}{576m^4} \right) \langle \frac{\alpha_s}{\pi} G^2 \rangle \right\} . \quad (51)$$

Within the full theory (SM) at quark level, the weak current is :

$$J_k^\alpha = \overline{q}_{kL} \gamma^\alpha Q \quad (52)$$

where Q is the heavy quark field in the full theory. Within HQEFT this current will, below the renormalization scale $\mu = m_Q (= m_b, m_c)$, be modified in the following way:

$$J_k^\alpha = \overline{\chi}_h \xi_{hk}^\dagger \Gamma^\alpha Q_v + \mathcal{O}(m_Q^{-1}) , \quad (53)$$




FIG. 7: Diagrams for bosonization of the left handed quark current.

The operator in equation (53) can be bosonized by calculating the Feynman diagrams shown in Fig. 7 which gives the bosonized current in (15) with:

$$\alpha_H \equiv -2iG_H N_c \left(-I_1 + mI_{3/2} + \frac{i(3\pi - 4)}{384N_c m^2} \langle \frac{\alpha_s}{\pi} G^2 \rangle \right). \quad (54)$$

To first order in the chiral expansion we obtain the bosonized current obtained by attaching one extra field $\mathcal{A}_\nu$ to the quark loops in Fig. 7:

$$J_k^\alpha(1) = \frac{1}{2} \text{Tr} \left[\xi_{hk}^\dagger \Gamma^\alpha H_{vh} (\alpha_{H\gamma}^{(1)} \gamma^\nu \gamma_5 + \alpha_{Hv}^{(1)} v^\nu \gamma_5) \mathcal{A}_\nu \right], \quad (55)$$

where the quantities $\alpha_H^{(1)}$ are given by expressions similar to (54).

The coupling α_H in (15) is related to the physical decay constants f_H and f_{H^*} , in the following way (for $H = B, D$):

$$\langle 0 | \bar{u} \gamma^\alpha \gamma_5 b | H \rangle = -2 \langle 0 | J_a^\alpha | H \rangle = iM_H f_H v^\alpha, \quad (56)$$

Taking the trace over the gamma matrices in (15), we obtain a relation for α_H and the relations between the heavy meson decay constants f_H and f_{H^*} (for $H = B, D$):

$$\alpha_H = \frac{f_H \sqrt{M_H}}{C_\gamma(\mu) + C_v(\mu)} = \frac{f_{H^*} \sqrt{M_{H^*}}}{C_\gamma(\mu)}, \quad (57)$$

where the model dictates us to put $\mu = \Lambda_\chi$. Later, in section V-B, we will see how chiral corrections and $1/m_Q$ corrections modify this relation.

C. Constraining the parameters of the HL χ QM

The gluon condensate can be related to the chromomagnetic interaction :

$$\mu_G^2(H) = \frac{1}{2M_H} C_M(\mu) \langle H | \bar{Q}_v \frac{1}{2} \sigma \cdot G Q_v | H \rangle, \quad (58)$$

where the coefficient $C_M(\mu)$ contains the short distance effects down to the scale μ and has been calculated to next to leading order (*NLO*) [26]. The chromomagnetic operator is responsible for the splitting between the 1^- and 0^- state, and is known from spectroscopy,

$$\mu_G^2(H) = \frac{3}{2}m_Q(M_{H^*} - M_H) . \quad (59)$$

An explicit calculation of the matrix element in equation (58) gives

$$\mu_G^2 = \eta_2 \frac{G_H^2}{m} \langle \frac{\alpha_s}{\pi} G^2 \rangle , \quad \text{where} \quad \eta_2 \equiv \frac{(\pi + 2)}{32} C_M(\Lambda_\chi) . \quad (60)$$

Combining the eqs. (31), (34) and (60) we obtain the following relations :

$$\langle \frac{\alpha_s}{\pi} G^2 \rangle = \frac{\mu_G^2 f^2}{2\eta_2} \frac{1}{\rho} , \quad G_H^2 = \frac{2m}{f^2} \rho , \quad (61)$$

where the quantity ρ is of order one and given by

$$\rho \equiv \frac{(1 + 3g_A) + \frac{\eta_1 \mu_G^2}{\eta_2 m^2}}{4(1 + \frac{N_c m^2}{8\pi f^2})} , \quad (62)$$

where $\eta_1 \equiv \frac{\pi}{32}$. In the limit where only the leading logarithmic integral I_2 is kept in (45), we obtain:

$$g_A \rightarrow 1 , \quad \rho \rightarrow 1 , \quad \beta \rightarrow \frac{1}{m} \quad G_H \rightarrow G_H^{(0)} \equiv \frac{\sqrt{2m}}{f} , \quad (63)$$

which for g_A and β correspond to the non-relativistic values.

From the eqs. (31), (50), and (60), we find

$$2\lambda_1 = \frac{1}{2}(3g_A - 1) + \frac{(9\pi - 16)\mu_G^2}{384\eta_2 m^2} . \quad (64)$$

In the limit (63) we obtain $2\lambda_1 \rightarrow 1$, as expected. The parameter λ_1 is related to the mass difference $M_{H_s} - M_{H_d}$. To leading order, we obtained the following expression for the β -term:

$$\beta = \frac{\rho}{m} \left\{ 1 + \frac{N_c m^2}{4\pi f^2} - \left(\frac{56 + 3\pi}{576 f^2 m^2} \right) \langle \frac{\alpha_s}{\pi} G^2 \rangle \right\} , \quad (65)$$

which is rather sensitive to m . Choosing m in the range 250-300 MeV we find [27] $\beta = (2.5 \pm 0.6) \text{ GeV}^{-1}$ to be compared with $\beta = (2.7 \pm 0.20) \text{ GeV}^{-1}$ extracted from experiment.

Using equation (45) and (34) we may write α_H as:

$$\alpha_H = \frac{G_H}{2} \left(-\frac{\langle \bar{q}q \rangle}{m} - 2f_\pi^2 \left(1 - \frac{1}{\rho} \right) + \frac{(\pi - 2)}{16m^2} \langle \frac{\alpha_s}{\pi} G^2 \rangle \right) , \quad (66)$$




FIG. 8: Loop diagrams for bosonizing the $b \rightarrow c$ current, $V^\mu = \gamma^\mu$, $A^\mu = \gamma^\mu \gamma_5$.

Combining (57) with (66), we obtain [12] in the leading limit (taking into account the logarithmic and quadratic divergent integrals only, and let $C_\gamma \rightarrow 1$, $C_v \rightarrow 0$ and $g_A \rightarrow 1$ as in (63)) we obtain the “Goldberger-Treiman like” relation:

$$f_H \sqrt{M_H} \rightarrow -\frac{\langle \bar{q}q \rangle}{f_\pi \sqrt{2m}} , \quad (67)$$

which gives the scale for f_H (It is, however, numerically a factor 2 off for the B -meson).

Using the relations (45), (47) and (62) we obtain for $\alpha_{H\gamma}^{(1)}$ and $\alpha_{Hv}^{(1)}$ in (55):

$$\alpha_{H\gamma}^{(1)} = \frac{2g_A}{G_H} , \quad (68)$$

$$\alpha_{Hv}^{(1)} = \frac{4}{3} G_H \left(\frac{f_\pi^2}{2m} \left(\frac{1}{\rho} - 1 \right) + \left[\frac{mN_c}{8\pi} + \frac{(\pi+8)}{256m^3} \langle \frac{\alpha_s}{\pi} G^2 \rangle \right] \right) . \quad (69)$$

Moreover, for the mass correction to the weak current given in (17) we find that $\omega_1 = -4\lambda_1/G_H$, where λ_1 is given in equation (50) or (64). The term ω'_1 is subleading in $1/N_c$.

The Isgur-Wise function $\zeta(\omega)$ in (19) relates all the form factors describing the processes $B(B^*) \rightarrow D(D^*)$ in the heavy quark limit. This function can be calculated from the diagrams shown in Fig. 8. The result before $1/m_Q$ and chiral corrections is:

$$\zeta(\omega) = \frac{2}{1+\omega} (1 - \rho) + \rho r(\omega) , \quad (70)$$

where ρ is given in (62) and $r(\omega)$ is the same function appearing in loop calculations of the anomalous dimension in HQEFT :

$$r(\omega) = \frac{1}{\sqrt{\omega^2 - 1}} \ln \left(\omega + \sqrt{\omega^2 - 1} \right) . \quad (71)$$

Note that $\zeta(1) = 1$ as it should.

D. The formal limit $m \rightarrow 0$

In this subsection we will discuss the limit of restoration of chiral symmetry, i.e. the limit $m \rightarrow 0$ [27]. In order to do this, we have to consider the various constraints obtained when constructing the HL χ QM [9].

Looking at the equations (31) and (34), one may worry that $\langle \bar{q}q \rangle$ and f behaves like $1/m$ in the limit $m \rightarrow 0$ unless one assumes that $\langle \frac{\alpha_s}{\pi} G^2 \rangle$ also go to zero in this limit. We should stress that the exact limit $m = 0$ cannot be taken because our loop integrals will then be meaningless. Still we may let m approach zero without going to this exact limit. In the pure light sector (at least when vector mesons are not included) there are no restrictions on how $\langle \frac{\alpha_s}{\pi} G^2 \rangle$ might go to zero. In the heavy light sector we have in addition to (31) and (34) also the relations (45) and (47), which put restrictions on the behavior of the gluon condensate $\langle \frac{\alpha_s}{\pi} G^2 \rangle$ for small masses. As $\langle \frac{\alpha_s}{\pi} G^2 \rangle$ has dimension mass to the fourth power, we find that $\langle \bar{q}q \rangle$ and f^2 may go to zero if $\langle \frac{\alpha_s}{\pi} G^2 \rangle$ goes to zero as m^4 or $m^3 \Lambda$ (eventually combined with $\ln(m/\Lambda)$). However, the behavior $m^3 \Lambda$ is inconsistent with the additional equations (61) and (62). Still, from all equations (45), (61), (62), we find the possible solution

$$\langle \frac{\alpha_s}{\pi} G^2 \rangle = \hat{c} N_c m^4 K(m) \quad , \quad \text{where} \quad K(m) \equiv (-4iI_2 + \frac{1}{8\pi}) \quad , \quad (72)$$

and $\hat{c}$ is some constant. Then we must have the following behavior for G_H^2 , g_A and μ_G^2 when m approaches zero:

$$G_H^2 \sim \frac{1}{N_c \Lambda} \quad , \quad (1 + 3g_A) \sim \frac{m}{\Lambda} K(m) \quad , \quad \mu_G^2 \sim \frac{m^3}{\Lambda} K(m) \quad , \quad (73)$$

with some restrictions on the proportionality factors. Here, the regularized I_2 is such that for small m , $K(m) = (c_1 + c_2 \ln m/\Lambda)$, c_1 and c_2 being constants. The behavior of G_H^2 is in agreement with Nambu-Jona-Lasinio models [1]. Note that in our model, $\delta g_A \rightarrow 4/3$ (corresponding to $g_A \rightarrow -1/3$) for $m \rightarrow 0$, in contrast to $\delta g_A \rightarrow 2/3$ for a free Dirac particle with $m = 0$. Note that in [9] we gave the variation of the gluon condensate with m for a fixed value of μ_G^2 . For the considerations in this subsection, we have to let μ_G^2 go to zero with m in order to be consistent. When $m \rightarrow 0$, we also find that $\beta \rightarrow 1/\Lambda$, provided that the coefficient $\hat{c}$ in (72) is fixed to a specific value (which is $\hat{c} = 576/(3\pi + 32) \simeq (1.93)^4$).




FIG. 9: Diagrams responsible for $1/m_Q$ terms in the chiral Lagrangian.

V. $1/m_Q$ CORRECTIONS WITHIN THE HL χ QM

A. Bosonization of the strong sector

To order $1/m_Q$ one obtains further contributions to chiral Lagrangians (see ref. [9] and references therein):

$$\begin{aligned}
\mathcal{L}_{Str} &= -\frac{\varepsilon_1}{m_Q} Tr [\overline{H}_k (i v \cdot D) H_k] + \frac{\varepsilon_1}{m_Q} Tr [\overline{H}_k H_h v_\mu \mathcal{V}_{hk}^\mu] \\
&+ \frac{g_1}{m_Q} Tr [\overline{H}_k H_h \gamma_\mu \gamma_5 \mathcal{A}_{hk}^\mu] + \frac{\varepsilon_2}{m_Q} Tr [\overline{H}_k \sigma^{\alpha\beta} i v \cdot D H_k \sigma_{\alpha\beta}] \\
&- \frac{\varepsilon_2}{m_Q} Tr [\overline{H}_k \sigma^{\alpha\beta} v_\mu \mathcal{V}_{hk}^\mu \sigma_{\alpha\beta} H_h] + \frac{g_2}{m_Q} Tr [\overline{H}_k \gamma_\mu \gamma_5 \mathcal{A}_{hk}^\mu H_h] + \dots
\end{aligned} \tag{74}$$

where the ellipses indicate other terms (of higher order, say), and D_μ contains the photon field. The new terms of order $1/m_Q$ in (74) are a consequence of the chromomagnetic interaction O_{mag} (the second term in equation (43)), and the kinetic interaction O_{kin} (the

third term in (43)). Calculating the diagrams of Fig. 9 and eliminating the divergent integrals and using equations (32), (34) and (48), gives for example

$$g_1 = m - G_H^2 \left(\frac{\langle \bar{q}q \rangle}{12m} + \frac{f_\pi^2}{6} + \frac{N_c m^2 (3\pi + 4)}{48\pi} - \frac{C_K}{16} \left(\frac{\langle \bar{q}q \rangle}{m} + 3f_\pi^2 \right) + \frac{(C_K - 2\pi)}{64m^2} \left\langle \frac{\alpha_s}{\pi} G^2 \right\rangle \right), \quad (75)$$

$$g_2 = \frac{(\pi + 4)}{(\pi + 2)} \frac{\mu_G^2}{6m}. \quad (76)$$

As the $1/m_Q$ terms break heavy quark spin symmetry, the chiral Lagrangian in (74) will split in $H(0^-)$ and $H^*(1^-)$ terms respectively.

B. The weak current to order $1/m_Q$

In HQEFT the weak vector current at order $1/m_Q$ is [8]:

$$J^\alpha = \sum_{i=1,2} C_i(\mu) J_i^\alpha + \frac{1}{2m_Q} \sum_j B_j(\mu) O_j^\alpha + \frac{1}{2m_Q} \sum_k A_k(\mu) T_k^\alpha, \quad (77)$$

where the first terms are given in (18) and (53), the B_j 's and A_j 's are Wilson coefficients, and the O_j^α 's are two quark operators

$$\begin{aligned} O_1^\alpha &= \bar{q}_L \gamma^\alpha i \not{D} Q_v, & O_4^\alpha &= \bar{q}_L \gamma^\alpha (-iv \cdot \overleftarrow{D}) Q_v, \\ O_2^\alpha &= \bar{q}_L v^\alpha i \not{D} Q_v, & O_5^\alpha &= \bar{q}_L v^\alpha (-iv \cdot \overleftarrow{D}) Q_v, \\ O_3^\alpha &= \bar{q}_L i D^\alpha Q_v, & O_6^\alpha &= \bar{q}_L (-i \overleftarrow{D}^\alpha) Q_v, \end{aligned} \quad (78)$$

The operators T_k are nonlocal and is a combination of the leading order currents J_i and a term of order $1/m_Q$ from the effective Lagrangian (43).

Combining (77) with (15), and adding chiral corrections and the $1/m_Q$ corrections indicated in (53), we obtain for $H = B, D$:

$$f_H = \frac{1}{\sqrt{M_H}} \left[(C_\gamma(\mu) + C_v(\mu)) \alpha_H + \frac{\eta_Q}{m_Q} + \frac{\eta_\chi}{32\pi^2 f_\pi^2} \right], \quad (79)$$

where $C_{\gamma,v}$ are defined in (18). Here the model dictates us to put $\mu = \Lambda_\chi$. The quantities η_Q and η_χ are given in [9]. One should note that the quantities η_Q for $Q = b, c$ depend on the Wilson coefficients C_i, B_i, A_i in (77) and some hadronic parameters, for instance $\varepsilon_{1,2}$ from (74). The Wilson coefficients entering f_H depends on m_Q through $\ln(m_Q/\mu)$, and therefore




FIG. 10: f_B as a function of m for $\langle \bar{q}q \rangle^{1/3} = -240$ MeV.

f_H is a complicated function of m_Q , $\langle \bar{q}q \rangle$, g_A , f_π , and the constituent light quark mass m . Note that $\langle \frac{\alpha_s}{\pi} G^2 \rangle^{1/4}$ is fixed to be around 320 MeV. In Fig. 10, f_B is plotted as function of m for standard values of the other parameters. One should note that bigger values of $|\langle \bar{q}q \rangle|$ give higher values of f_B . For a discussion of the numerical values of our parameters, see [9] and [27].

VI. APPLICATIONS

In this section we focus on the chiral quark model aspects, especially contributions proportional to the gluon condensate. There are always additional chiral loop corrections which can be found in [5, 9, 13, 14] and references therein.

A. $B - \bar{B}$ mixing and heavy quark effective theory

At quark level, the standard effective Lagrangian describing $B - \bar{B}$ mixing is [28]

$$\mathcal{L}_{eff}^{\Delta B=2} = -\frac{G_F^2}{4\pi^2} M_W^2 (V_{tb}^* V_{tq})^2 S_{IL}(x_t) \eta_B b(\mu) Q_B, \quad (80)$$

where G_F is Fermi's coupling constant, the V 's are KM factors (for which $q = d$ or s for B_d and B_s respectively) and S_{IL} is the Inami-Lim function due to short distance electroweak

loop effects for the box diagram. The quantity $Q_B \equiv Q(\Delta B = 2)$ is a four quark operator

$$Q_B = \overline{q}_L \gamma^\alpha b_L \overline{q}_L \gamma_\alpha b_L , \quad (81)$$

where q_L (b_L) is the left-handed projection of the q (b)-quark field. The quantities $\eta_B = 0.55 \pm 0.01$ and $b(\mu)$ are calculated in perturbative quantum chromodynamics (QCD). At the renormalization scale $\mu = m_b$ ($\simeq 4.8$ GeV) one has $b(m_b) \simeq 1.56$ in the naive dimension regularization scheme. The matrix element of the operator Q_B between the meson states is parameterized by the bag parameter B_{B_q} :

$$\langle B | Q_B | \overline{B} \rangle \equiv \frac{2}{3} f_B^2 M_B^2 B_{B_q}(\mu) , \quad (82)$$

where by definition, $B_{B_q} = 1$ within the factorized limit. In general, the matrix element of the operator Q_B is dependent on μ , and thereby B_{B_q} also depends on μ . As for $K - \overline{K}$ mixing, one defines a renormalization scale independent quantity

$$\hat{B}_{B_q} \equiv b(\mu) B_{B_q}(\mu) . \quad (83)$$

The $\Delta B = 2$ operator in equation (81) can for $\Lambda_\chi < \mu < m_b$ be written [29, 30] :

$$Q_B = C_1 Q_1 + C_2 Q_2 + \frac{1}{m_b} \sum_i h_i X_i + \mathcal{O}(1/m_b^2) . \quad (84)$$

The operator Q_1 is Q_B for b replaced by $Q_v^{(\pm)}$, while Q_2 is generated within perturbative QCD for $\mu < m_b$. The operators X_i are taking care of $1/m_b$ corrections. The quantities C_1, C_2, h_i are Wilson coefficients. The operators are given by

$$Q_1 = 2 \overline{q}_L \gamma^\mu Q_v^{(+)} \overline{q}_L \gamma_\mu Q_v^{(-)} , \quad (85)$$

$$Q_2 = 2 \overline{q}_L v^\mu Q_v^{(+)} \overline{q}_L v_\mu Q_v^{(-)} , \quad (86)$$

$$X_1 = 2 \overline{q}_L i D^\mu Q_v^{(+)} \overline{q}_L \gamma_\mu Q_v^{(-)} + \dots \quad (87)$$

The explicit expressions for the operators X_i are given in [13]. There are also non-local operators constructed as time-ordered products of $Q_{1,2}$ and the first order HQEFT Lagrangian in (43). The Wilson coefficients C_1 and C_2 have been calculated to NLO [29] and for $\mu = \Lambda_\chi$ one has $C_1(\Lambda_\chi) = 1.22$ and $C_2(\Lambda_\chi) = -0.15$. The coefficients h_i have been calculated to leading order (LO) in [30].




FIG. 11: Non-factorizable soft gluonic contribution to the bag-parameter. (Here $\Gamma \equiv t^a \gamma^\mu L$.)

In order to find all the matrix element of $Q_{1,2}$, one uses the following relation between the generators of $SU(3)_c$ (i, j, l, n are colour indices running from 1 to 3):

$$\delta_{ij}\delta_{ln} = \frac{1}{N_c}\delta_{in}\delta_{lj} + 2 t_{in}^a t_{lj}^a, \quad (88)$$

where a is an index running over the eight gluon charges. This means that by means of a Fierz transformation, the operator Q_1 in (85) may also be written in the following way (there is a similar expression for Q_2):

$$Q_1^F = \frac{1}{N_c}Q_1 + 4 \bar{q}_L t^a \gamma^\mu Q_v^{(+)} \bar{q}_L t^a \gamma_\mu Q_v^{(-)}. \quad (89)$$

The first (naive) step to calculate the matrix element of a four quark operator like Q_1 is to insert vacuum states between the two currents. This factorized limit means to bosonize the two currents in Q_1 and multiply them (see (53)). The second operator in (89) is genuinely non-factorizable. In the approximation where only the lowest gluon condensate is taken into account, the last term in (89) can be written in a *quasi-factorizable* way by bosonizing the heavy-light colour current with an extra colour matrix t^a inserted and with an extra gluon emitted as shown in Fig. 11.

We find the bosonized colour current:

$$(\bar{q}_L t^a \gamma^\alpha Q_v^{(\pm)})_{1G} \longrightarrow -\frac{G_H g_s}{8} G_{\mu\nu}^a \text{Tr} \left[\xi^\dagger \gamma^\alpha L H^{(\pm)} \left(\pm i I_2 \{ \sigma^{\mu\nu}, \gamma \cdot v \} + \frac{1}{8\pi} \sigma^{\mu\nu} \right) \right], \quad (90)$$

where $\{, \}$ symbolizes an anti-commutator. The result for the right part of the diagram with $\bar{B}$ replaced by B is obtained by changing the sign of v and letting $P_5^{(+)} \rightarrow P_5^{(-)}$. Multiplying the coloured currents, we obtain the non-factorizable parts of Q_1 and Q_2 to first order in the gluon condensate by using eq. (30).

Now the bag parameter can be extracted and may be written in the form:

$$\hat{B}_{B_q} = \frac{3}{4} \tilde{b} \left[1 + \frac{1}{N_c} (1 - \delta_G^B) + \frac{\tau_b}{m_b} + \frac{\tau_\chi}{32\pi^2 f^2} \right], \quad (91)$$




FIG. 12: The bag parameter $\hat{B}$ for B_s as a function of m .

where the parameter $\tilde{b}$ also involves the Wilson coefficients $C_{\gamma,v}$ defined in (18):

$$\tilde{b} = b(m_b) \left[\frac{C_1 - C_2}{(C_\gamma + C_v)^2} \right]_{\mu=\Lambda_\chi}. \quad (92)$$

The soft gluonic non-factorizable effects are given by

$$\delta_G^B = \frac{N_c \langle \frac{\alpha_s}{\pi} G^2 \rangle}{32\pi^2 f^2 f_B^2} \frac{m}{M_B} \kappa_B \left[\frac{C_1}{C_1 - C_2} \right]_{\mu=\Lambda_\chi}, \quad (93)$$

where κ_B is a dimensionless hadronic parameter which depends on m, f, μ_G^2 and g_A and is of order 2. Note that we are qualitatively in agreement with [31], where also a negative contribution to the bag factor from soft gluon effects is found. Numerically, f and f_B are of the same order of magnitude, and δ_G^B is therefore suppressed like m/M_B compared to the corresponding quantity

$$\delta_G^K = N_c \frac{\langle \frac{\alpha_s}{\pi} G^2 \rangle}{32\pi^2 f^4} \quad (94)$$

found for $K - \bar{K}$ mixing [5].

However, one should note that f_B scales as $1/\sqrt{M_B}$ within HQEFT, and therefore δ_G^B is still formally of order $(m_b)^0$. The quantity τ_b represents the $1/m_b$ corrections due to the operators X_i . Furthermore, the quantity τ_χ represents the chiral corrections (including counterterms) to the bosonized versions of $Q_{1,2}$ [13]. The bag parameter $\hat{B}$ is plotted as function of m in Fig. 12 for the case B_s . Our results are numerically in agreement with lattice results [32].

B. The processes $B \rightarrow D^{(*)}\overline{D}^{(*)}$

It has been observed[14] that the processes $\bar{B}_{d,s}^0 \rightarrow D_{s,d}\bar{D}_{s,d}$, $\bar{B}_{d,s}^0 \rightarrow D_{s,d}^*\bar{D}_{s,d}$, $\bar{B}_{d,s}^0 \rightarrow D_{s,d}\bar{D}_{s,d}^*$, and $\bar{B}_{d,s}^0 \rightarrow D_{s,d}^*\bar{D}_{s,d}^*$, have no factorized contribution from the spectator mechanism. If one or two of the D -mesons in the final state are vectors, there are relatively small contributions from the annihilation mechanism. The effective non-leptonic Lagrangian at quark level has the usual form [28]:

$$\mathcal{L}_W = -4\frac{G_F}{\sqrt{2}}V_{cb}V_{cq}^* \sum_i a_i \hat{Q}_i(\mu). \quad (95)$$

In our case there are only two numerically relevant operators (for $q = d, s$):

$$\hat{Q}_1 = (\bar{q}_L \gamma^\alpha b_L) (\bar{c}_L \gamma_\alpha c_L) ; \hat{Q}_2 = (\bar{c}_L \gamma^\alpha b_L) (\bar{q}_L \gamma_\alpha c_L). \quad (96)$$

At $\mu = m_b$, one has $a_2 \sim 1$ and $a_1 \sim 1/10$.

Using (88), we obtain the Fierzed version of the operators $\hat{Q}_{1,2}$:

$$\begin{aligned} \hat{Q}_1^F &= \frac{1}{N_c} \hat{Q}_2 + 2(\bar{c}_L \gamma^\alpha t^a b_L) (\bar{q}_L \gamma_\alpha t^a c_L) \\ \hat{Q}_2^F &= \frac{1}{N_c} \hat{Q}_1 + 2(\bar{q}_L \gamma^\alpha t^a b_L) (\bar{c}_L \gamma_\alpha t^a c_L) \end{aligned} \quad (97)$$

The genuine non-factorizable $1/N_c$ chiral Lagrangian terms from “coloured quark operators” can be estimated within the $HL\chi QM$. However, in order to do this we have to treat the effective weak non-leptonic Lagrangian in (95) within heavy quark effective theory (HQEFT) [8]. Then b , c , and $\bar{c}$ quarks are replaced by their corresponding operators in HQEFT:

$$b \rightarrow Q_{v_b}^{(+)} , \quad c \rightarrow Q_{v_c}^{(+)} , \quad \bar{c} \rightarrow Q_{\bar{v}}^{(-)} \quad (98)$$

up to $1/m_b$ and $1/m_c$ corrections. Then the effective weak non-leptonic Lagrangian (95) can be evolved down to the scale $\mu \sim \Lambda_\chi \sim 1$ GeV [33]. At $\mu = 1$ GeV we have $a_2 \simeq 1.29 + 0.08i$, and $a_1 \simeq -0.35 - 0.07i$. Note that $a_{1,2}$ are complex for $\Lambda_\chi < \mu < m_c$ [33].

The bosonized factorized weak Lagrangian corresponding to Fig. 13 and the operator $\hat{Q}_2$ (with the dominating Wilson coefficient a_2) is obtained from (15), and (19), and (95):

$$\mathcal{L}_{W-Fact}^{Bos}(Q_2) = 4\frac{G_F}{\sqrt{2}}V_{cb}V_{cq}^* (a_2 + \frac{a_1}{N_c}) \zeta(\omega) \frac{\alpha_H}{2} Tr \left[\overline{H_c^{(+)}} \gamma^\alpha L H_b^{(+)} \right] \cdot Tr \left[\xi^\dagger \gamma^\alpha L H_{\bar{c}}^{(-)} \right] , \quad (99)$$

where $\omega \equiv v_b \cdot v_c = v_b \cdot \bar{v} = M_B/(2M_D)$. This Lagrangian (corresponding to the spectator mechanism) contributes to the factorized amplitude for the process $\bar{B}^0 \rightarrow D^+ D_s^-$, and is the




FIG. 13: Factorized contribution for $\overline{B}^0 \rightarrow D^+ D_s^-$ through the spectator mechanism, which does not exist for the decay mode $\overline{B}^0 \rightarrow D_s^+ D_s^-$. The double dashed lines represent heavy mesons, the double lines represent heavy quarks, and the single lines light quarks.




FIG. 14: Factorized contribution for $\overline{B}^0 \rightarrow D_s^+ D_s^-$ through the annihilation mechanism, which give zero contributions if both D_s^+ and D_s^- are pseudo-scalars.

starting point for chiral loop contributions of order $(m_K g_A / 4\pi f)^2$ (which are $1/N_c$ suppressed) to the processes $B \rightarrow D^{(*)} \overline{D}^{(*)}$.

The bosonized factorized weak Lagrangian corresponding to Fig. 14 and the non-dominating Wilson coefficient a_1 is

$$\mathcal{L}_{W-Fact}^{Bos}(Q_1) = 4 \frac{G_F}{\sqrt{2}} V_{cb} V_{cq}^* (a_1 + \frac{a_2}{N_c}) \zeta(-\lambda) \frac{\alpha_H}{2} Tr \left[\xi^\dagger \gamma^\mu L H_b^{(+)} \right] \cdot Tr \left[\overline{H_c^{(+)}} \gamma^\alpha L H_c^{(-)} \right] \quad (100)$$

where $\lambda \equiv \bar{v} \cdot v_c = (M_B^2 / (2M_D^2) - 1)$. Unless one or both of the D -mesons in the final state are vector mesons, this matrix element is zero due to current conservation, which is analogous to the decay mode $\overline{D}^0 \rightarrow K^0 \overline{K}^0$ [12].

The genuine non-factorizable part for $\overline{B}^0 \rightarrow D_s^+ D_s^-$ at quark level can, by means of Fierz transformations and the identity (88), be written in terms of colour currents. The left part in Fig. 15 with gluon emission gives us the bosonized colour current which is the same as




FIG. 15: Non-factorizable contribution for $\overline{B}^0 \rightarrow D_s^+ D_s^-$ through the annihilation mechanism with additional soft gluon emission. The wavy lines represent soft gluons ending in vacuum to make gluon condensates.

for $B - \overline{B}$ mixing in eq. (90). For the creation of a $D\overline{D}$ pair in the right part of Fig. 15, there is an analogue of (90), which can be written:

$$\left(\overline{Q_{v_c}^{(+)}} t^a \gamma^\alpha L Q_{\bar{v}}^{(-)} \right)_{1G} \longrightarrow \frac{G_H^2 g_s}{32\pi m} G_{\mu\nu}^a \text{Tr} \left[\overline{H_c^{(+)}} \gamma^\alpha L H_{\bar{c}}^{(-)} X^{\mu\nu} \right] \quad (101)$$

where

$$X^{\mu\nu} \equiv \frac{r(-\lambda)}{\pi} \sigma^{\mu\nu} + \frac{1}{4(\lambda-1)} \{ \sigma^{\mu\nu}, \gamma \cdot t \}$$

and $t \equiv v_c - \bar{v}$. Multiplying the currents and using (30) we obtain a bosonized effective Lagrangian as the product of two traces. Note that our non-factorizable amplitudes (proportional to the gluon condensate) are proportional to the numerically favourable Wilson coefficient a_2 .


The gluon condensate contribution obtained from (90) and (101) is a linear combination of terms like:

$$\text{Tr} \left[\xi^\dagger \sigma^{\mu\alpha} L H_b^{(+)} \right] \cdot \text{Tr} \left[\overline{H_c^{(+)}} \gamma_\alpha L H_{\bar{c}}^{(-)} \gamma_\mu \right] ,$$

$$\text{Tr} \left[\xi^\dagger \gamma^\mu L H_b^{(+)} \right] \cdot \text{Tr} \left[\overline{H_c^{(+)}} \gamma^\alpha L H_{\bar{c}}^{(-)} \sigma_{\mu\alpha} R \right] ,$$

$$\text{Tr} \left[\xi^\dagger \gamma^\mu L H_b^{(+)} \right] \cdot \text{Tr} \left[\overline{H_c^{(+)}} \gamma_5 L H_{\bar{c}}^{(-)} \gamma_\mu \right] ,$$

$$\text{Tr} \left[\xi^\dagger L H_b^{(+)} \right] \cdot \text{Tr} \left[\overline{H_c^{(+)}} \gamma^\alpha L H_{\bar{c}}^{(-)} \gamma_\alpha \right] ,$$

FIG. 16: Gluon condensate contributions to $B \rightarrow D\eta'$.

$$Tr \left[\xi^\dagger \sigma^{\mu\alpha} L H_b^{(+)} \right] \cdot Tr \left[\overline{H_c^{(+)}} \gamma_\alpha L H_{\bar{c}}^{(-)} \right] (\bar{v} - v_c)_\mu ,$$

$$\varepsilon^{\mu\nu\alpha\lambda} (v_c + \bar{v})_\nu Tr \left[\xi^\dagger \gamma^\mu L H_b^{(+)} \right] \cdot Tr \left[\overline{H_c^{(+)}} \gamma^\alpha L H_{\bar{c}}^{(-)} \gamma_\lambda \right] . \quad (102)$$

These terms might have been written down based on the heavy quark symmetry, but the HL χ QM selects a certain linear combination to be realized.

Our amplitudes for $B \rightarrow D\bar{D}$, in terms of chiral loop and gluon condensate contributions, are sensitive to $1/m_c$ corrections and counterterms which are not yet calculated [14]. Operators suppressed by $1/m_Q$ are obtained by the replacements

$$Q_v^{(\pm)} \rightarrow \frac{1}{m_Q} i\gamma \cdot D_\perp Q_v^{(\pm)} \quad ; \quad D_\perp^\nu = D^\nu - v^\nu (v \cdot D) , \quad (103)$$

for one of the heavy quarks in (96) and (97). Some new quark operators of order $1/m_Q$ might also be generated by pQCD for $\mu < m_Q$. Counterterms correspond to mass insertion of $\widetilde{M}_q$, given by (14) and (26), at light quark lines in the diagrams for $B \rightarrow D\bar{D}$ this subsection.

C. Other applications

Within the HL χ QM, the process $B \rightarrow D\eta'$ has been estimated [11]. This is done in two steps. First we calculate the subprocess $B \rightarrow Dgg^*$. Then the virtual gluon g^* is attached to the $\eta'gg^*$ -vertex, and the other end in vacuum and make a gluon condensate together with one of the other soft gluons (g) from the $\eta'gg^*$ -vertex. Using Fierz transformations for the four quark operators for $b \rightarrow cd\bar{u}$, we obtain contributions corresponding to Fig. 16.

We have used existing parameterizations of the $\eta'gg^*$ -vertex form factor and assumed that the current for $B \rightarrow g^*$ is related to the better known case $B \rightarrow \rho$. It turns out that the “factorizable” diagram to the left in figure 16 can be neglected compared to the non-factorizable diagram to the right. For m in the range 230-270 MeV, we obtained the

result [11] $Br(B \rightarrow D\eta') = (2.2 \pm 0.4) \times 10^{-4}$. Here $1/m_Q$ and chiral corrections are not included.

Heavy to light non-leptonic processes like $B, D \rightarrow K\pi$ cannot in general be treated within the HL χ QM in its present form. (See, however, section VII-B). Still, semileptonic heavy to light processes might be treated at the “no recoil point” [9]. The form factors $f_+(q^2)$ and $f_-(q^2)$ are defined as:

$$\langle \pi^+(p_\pi) | \bar{u}\gamma^\alpha b | H \rangle = 2 \langle \pi^+(p_\pi) | J_f^\alpha | H \rangle = f_+(q^2)(p_H + p_\pi)^\alpha + f_-(q^2)(p_H - p_\pi)^\alpha \quad (104)$$

where $p_H^\alpha = M_H v^\alpha$ and the index a corresponds to quark flavour u and $q^\mu = p_H^\mu - k_\pi^\mu$. The form factors get contributions from $J_f^\alpha(0)$ in (15) and $J_f^\alpha(1)$ in (55) close to the “no recoil point” where $v \cdot p_\pi$ is small:

$$f_+(q^2) + f_-(q^2) = \frac{-1}{\sqrt{2}M_H f_\pi} (C_\gamma + C_v - g_A C_\gamma) \alpha_H, \quad (105)$$

$$f_+(q^2) - f_-(q^2) = -C_\gamma \frac{\sqrt{M_H}}{\sqrt{2}f_\pi} \left(\frac{g_A \alpha_H}{v \cdot p_\pi} + \alpha_{H\gamma}^{(1)} \right), \quad (106)$$

where we have neglected terms of first order in $v \cdot p_\pi$ (where $\alpha_{Hv}^{(1)}$ contributes). The $1/v \cdot p_\pi$ term in (106) is due to the H^* pole. From equation (105) and (106) we see that

$$(f_+(q^2) + f_-(q^2))/(f_+(q^2) - f_-(q^2)) \sim 1/M_H \quad (107)$$

which is the well known Isgur-Wise scaling law [34]. The equations for the two form factors f_+ and f_- should be studied further, and chiral corrections and $1/m_Q$ corrections should be added.

VII. FURTHER POSSIBLE EXTENSIONS OF CHIRAL QUARK MODELS

In this section we consider two possible extensions of chiral quark models which are not yet worked out in detail. The descriptions are therefore sketchy.

A. Inclusion of light vectors

One might include vectors in the chiral perturbation theory [35] and thus it should be possible to use the chiral quark model also in this case. We suggest a Lagrangian

$$\mathcal{L} = \mathcal{L}_{mass} + \mathcal{L}_{\chi QM} + \mathcal{L}_{IVA}, \quad (108)$$

where the interaction between quarks and the vectors and axial vectors is given by

$$\mathcal{L}_{IVA} = \bar{\chi} [h_V \gamma^\mu V_\mu + h_A \gamma_5 A_\mu] \chi . \quad (109)$$

Here V are given as Π in (3) with π replaced by ρ etc, and similarly for the axial vector A where π is replaced by a_1 . The (bare) mass term is

$$\mathcal{L}_{mass} = \bar{m}_V^2 Tr [V_\mu V^\mu] + \bar{m}_A^2 Tr [A_\mu A^\mu] . \quad (110)$$

After quarks are integrated out, the masses are modified and identified with the physical ones. Then a kinetic term is also generated:

$$\mathcal{L}_{kin} = -\frac{1}{2} [V_{\mu\nu} V^{\mu\nu}] - \frac{1}{2} [A_{\mu\nu} A^{\mu\nu}] ; \quad (111)$$

where for $X = V, A$:

$$X_{\mu\nu} = \nabla_\mu X_\nu - \nabla_\nu X_\mu , \quad (112)$$

and similar for the axial vector. Here ∇ is a covariant derivative including the goldstones:

$$\nabla_\mu X_\nu \equiv \partial_\mu X_\nu + i [\mathcal{V}_\mu, X_\nu] . \quad (113)$$

For the left-handed current for $vac \rightarrow X = V, A$ we find the $SU(3)$ octet current

$$J_\mu^n(vac \rightarrow X) = \frac{1}{2} k_X Tr [\Lambda^n X_\mu] , \quad (114)$$

where the quantity Λ^n is given by (36).

As previously, bosonization gives constraints on the parameters of the vectorial sector. From normalization of the kinetic term(s) we obtain:

$$\frac{f^2 h_V^2}{3m^2} \left[1 - \frac{1}{15m^2 f^2} \langle \frac{\alpha_s}{\pi} G^2 \rangle \right] = 1 , \quad (115)$$

where $h_A = h_V$ before chiral corrections. For the currents we obtain

$$k_V = \frac{1}{2} h_V \left(-\frac{\langle \bar{q}q \rangle}{m} + f^2 - \frac{1}{8m^2} \langle \frac{\alpha_s}{\pi} G^2 \rangle \right) , \quad (116)$$

and similarly , for the axial case:

$$k_A = \frac{1}{2} h_A \left(-\frac{\langle \bar{q}q \rangle}{m} - 3f^2 + \frac{1}{8m^2} \langle \frac{\alpha_s}{\pi} G^2 \rangle \right) . \quad (117)$$

The formalism suggested in this subsection might for instance, when combined with HL χ QM, give a reasonable description of the weak current for D -meson decays $D \rightarrow V$ [36]. This might also be the case for processes like $D \rightarrow VP$, where V is a vector meson and P is a pseudoscalar. In the last case, non-factorizable contributions can be calculated in terms of chiral loops and gluon condensates. However, one should keep in mind that a limitation in this case is that V is (also formally) light compared to D .

B. Heavy to light transitions

As emphasized in the introduction, the HL χ QM is not suited to describe $B \rightarrow \pi$ transitions except for semileptonic transitions close to the no recoil point. It might therefore be surprising that we consider a formalism for chiral perturbation theory for $B \rightarrow \pi$ transitions (and more general $B \rightarrow P$ for $P = \pi, K, \eta$), because the involved pion is hard. However, in general, in a transition $B \rightarrow$ pions, we might have a configuration where one pion is hard and one (or more) is soft. For such cases we split the pseudoscalar sector in hard and soft pseudoscalars. The soft pseudoscalars are represented as before, while the hard pseudoscalars are represented by an octet 3×3 matrix M given as Π in eq. (3), but transforming as Σ under $SU(3)_L \times SU(3)_R$.

Starting with a γ_5 coupling for quarks coupling to pseudoscalars, we represent the hard light quark with a quark field q_n [16, 37] and the soft light quarks by the flavour rotated fields χ in section III. Then we arrive at an interaction Lagrangian

$$\mathcal{L}_n = G_M \bar{\chi} [\xi^\dagger M R - \xi M^\dagger L] q_n, \quad (118)$$

for a hard light quark q_n entering a hard pion (kaon) with momentum $p_M = E n$ where n is a lightlike vector and E is the energy of the hard pion(kaon). The hard quark has then momentum $p_q = E n + k$, where k is of order $\Lambda_\chi \sim 1$ GeV or smaller. G_M is a coupling which has to be determined by some physical requirements. For an outgoing hard quark we have

$$\mathcal{L}_{\bar{n}} = G_M \bar{q}_n [M \xi^\dagger R - M^\dagger \xi L] \chi. \quad (119)$$

Now one might combine (118) and (119) with HL χ QM and use some version of a large energy effective field theory (LEET) [37] to describe the light hard quarks. Using the LEET propagator

$$\frac{\gamma \cdot n}{2n \cdot k} \quad (120)$$

for the light hard quark, we can write down a quark loop diagram for $B \rightarrow P$ with a corresponding amplitude for the heavy-light weak current (to leading order)

$$J_k^\alpha(B \rightarrow P) = K \text{Tr} [\Gamma^\alpha H_{vh} \gamma \cdot n \xi M^\dagger] . \quad (121)$$

Given the transformation properties in (7), (12), and (24), the current (121) transforms as in (16).

The behaviour of the quantity (form factor) K is known from theoretical considerations within LEET[37] and soft collinear effective theory (SCET)[16]:

$$K \sim E \zeta^{(v)}(M_B, E) , \quad (122)$$

where $\zeta^{(v)}$ is expected to scale as

$$\zeta^{(v)}(M_B, E) \sim \frac{\sqrt{M_B}}{E^2} . \quad (123)$$

Note that a factor $\sqrt{M_B}$ is associated with the heavy (B) meson wave function and similarly a factor $\sqrt{E}$ with the wave function of the hard pseudoscalar meson. Within our framework, K will contain the product of the couplings G_H and G_M , and some loop integrals involving the heavy quark propagator, the ordinary Dirac propagator for the soft quark, and the LEET propagator in (120). However, it has been pointed out that the LEET propagator is too singular to give meaningful loop integrals[38], and that the LEET is incomplete[16]. Therefore the simple expression in (120) has to be modified in some way, by keeping $n^\mu n_\mu = \delta^2 \neq 0$ with $\delta \sim 1/E$, by adding a small quantity in the LEET propagator denominator, or by modifying the formalism in other ways. And this modification has to be done such that one does not come in conflict with the known scaling properties of $\zeta^{(v)}$. Keeping $\delta \neq 0$ and $\delta \sim 1/E$, some of the involved loop integrals have the same mathematical form as those involved in the Isgur-Wise function in (70), but with $\omega \rightarrow 1/\delta$. The most plausible scenario is that $G_M \sim E^{-3/2}$. Anyway, knowledge of $\zeta^{(v)}$ will put restrictions on G_M .

The $W \rightarrow \pi$ transition is in [15, 16] represented by an integral over a momentum distribution proportional to $x(1-x)$ dominated at $x \sim 1/2$. However, there are also suppressed contributions (for $E \gg \Lambda_\chi$) from momentum configurations where one quark (anti-quark) is hard and the anti-quark (quark) is soft. The left-handed current is in this case given by

$$j_\mu^l = \bar{q}_n \gamma_\mu L (\lambda^l \xi) \chi \quad , \quad \text{or} \quad j_\mu^l = \bar{\chi} (\xi^\dagger \lambda^l) \gamma_\mu L q_n , \quad (124)$$

where l is an $SU(3)$ octet index. These quark currents will, when combined with (118) and (119), generate a bosonized current

$$\Delta J_\mu^l = N \tilde{n}^\mu Tr [\lambda^l (\Sigma M^\dagger + M \Sigma^\dagger)] , \quad (125)$$

where $\tilde{n}$ is another (almost) lightlike vector with opposite three momentum compared to n such that $\tilde{n}^2 = \delta^2$ and $\tilde{n} \cdot n = 2 - \delta^2$. In the most plausible scenario mentioned above N scales as a constant (E^0), which is suppressed by $1/E$ compared with the leading order current $\sim E f_P^{(0)} \tilde{n}^\mu$. The physical decay constant f_P (for $P = \pi, K, \eta$) is within this scheme given by $f_P^{(0)}$ plus the suppressed contribution $\sim N/E$ from (125).

Now, the product of the currents in (121) and (125) will give a factorized $1/E$ suppressed contribution to $B \rightarrow K\pi$ corresponding to the diagram in Fig. 13, with $\bar{c}$ and c replaced by energetic (anti) quarks, D by π and D_s by K . This contribution can of course not be distinguished from the standard factorized contribution. However, pulling out soft pseudoscalars from ξ and Σ in the currents (121) and (125), we obtain $1/E$ suppressed non-factorizable chiral loop contributions to $B \rightarrow K\pi$. Similarly there will be $1/E$ suppressed gluon condensate contributions. Such suppressed terms are not in conflict with QCD factorization [15].

VIII. CONCLUSION

We have presented the main features of chiral quark models, both in the pure light and the heavy-light sector. Especially, the HL χ QM seem to work well. In that case, it is possible to systematically calculate the $1/m_Q$ corrections as well as chiral corrections. The model may be used to give predictions for many quantities. Especially, it is suitable for calculation of the B -parameter for $B - \bar{B}$ mixing [9], and for a study of processes of the type $B \rightarrow D\bar{D}$. For heavy to light transitions ($B \rightarrow K\pi$, say) the HL χ QM cannot be used in its present form. It remains to be seen if the extension indicated in sect VII-B to incorporate light energetic quarks will lead to some understanding of such decays.

In our version of the chiral quark models (pure light and heavy light cases) soft gluon effects are truncated to include only the second order gluon condensate. It has worked reasonably well up to now, but one may wonder if this is enough to accommodate all effects, for instance when light vectors are included. Maybe for instance higher order gluon condensates could be included, but our simple model will of course then be much more complicated.

Acknowledgments

J.O.E. is supported in part by the Norwegian research council and by the European Union RTN network, Contract No. HPRN-CT-2002-00311 (EURIDICE). He thanks S. Bertolini, M. Fabbrichesi, S. Fajfer, I. Picek and A. Polosa for collaboration and discussions on chiral quark models.

-
- [1] J. Bijnes, *Phys. Rept.* **265** (1996).
 - [2] J.A. Cronin, *Phys. Rev.* **161** (1967) 1483,
 S. Weinberg, *Physica* **96A** (1979) 327,
 Ebert and Volkov, *Z. Phys. C* **16**(1983) 205,
 A. Manohar and H. Georgi, *Nucl. Phys.* **B234**(1984) 189,
 J. Bijnens, H. Sonoda and M.B. Wise, *Can. J. Phys.* **64** (1986) 1,
 Ebert and Reinhardt, *Nucl. Phys.* **B71**(1986) 188,
 D.I. Diakonov, V.Yu. Petrov and P.V. Pobylitsa, *Nucl. Phys.* **B306**(1988) 809.
 - [3] J. O. Eeg and I. Picek, *Phys. Lett.* **B301** (1993) 423,
 J. O. Eeg and I. Picek, *Phys. Lett.* **B323** (1994) 193,
 A.E. Bergan and J.O. Eeg, *Phys. Lett.* **390** (1997) 420.
 - [4] D. Espriu, E. de Rafael and J. Taron, *Nucl. Phys.* **B345** (1990) 22. A. Pich and E. de Rafael, *Nucl. Phys.* **B358** (1991) 311. D. Ebert and M.K. Volkov, *Phys.Lett. B* **272** (1991), 86.
 - [5] S. Bertolini, J.O. Eeg and M. Fabbrichesi, *Nucl. Phys.* **B449** (1995) 197. V. Antonelli, S. Bertolini, J.O. Eeg, M. Fabbrichesi and E.I. Lashin, *Nucl. Phys.* **B469** (1996) 143. S. Bertolini, J.O. Eeg, M. Fabbrichesi and E.I. Lashin, *Nucl. Phys.* **B514** (1998) 63-92. and *ibid.* **B514** (1998) 93-112.
 - [6] M.A. Nowak, M. Rho, and I. Zahed, *Phys.Rev.* **D48** (1993) 4370.
 W.A. Bardeen and C.T. Hill, *Phys. Rev.* **D49** (1994) 409 .
 D. Ebert, T. Feldmann R. Friedrich and H. Reinhardt, *Nucl. Phys.* **B434** (1995) 619.
 A. Deandrea, N. Di Bartolomeo, R. Gatto, G. Nardulli, and A.D. Polosa, *Phys. Rev.* **D58** (1998) 034004. A. Polosa, hep - ph/0004183.
 D. Ebert, T. Feldmann and H. Reinhardt, *Phys.Lett.* **B388** (1996) 154.

- [7] M. Wise, *Phys. Rev. D* **45**, (1992) 2188 .
R. Casalbuoni, A. Deandrea, N. Di Bartolomeo, R. Gatto, F. Feruglio and G. Nardulli, *Phys. Rep.* **281** (1997) 145.
- [8] See for example: A.V. Manohar and M.B. Wise, in “Heavy Quark Physics”, published in Cambridge Monogr. Part. Phys. Nucl. Phys. Cosmol. 10 (2000),
M. Neubert, *Phys. Rep.* **245** (1994) 259.
- [9] A. Hiorth and J. O. Eeg, *Phys. Rev. D* **66** (2002) 074001.
- [10] H. Y. Cheng, *Chin. J. Phys.* **38** (2000) 1044, hep-ph/9911202.
- [11] J. O. Eeg, A. Hiorth, A. D. Polosa, *Phys. Rev. D* **65** (2002) 054030.
- [12] J. O. Eeg, S. Fajfer, and J. Zupan, *Phys. Rev. D* **64** (2001) 034010.
- [13] A. Hiorth and J. O. Eeg, *Eur.Phys.J.direct* **C30** (2003) 006.
- [14] Jan.O. Eeg, Svjetlana Fajfer, and Aksel Hiorth, *Phys. Lett. B* **570** (2003) 46-52).
J.O. Eeg, S. Fajfer, A. Hiorth, and A. Prapotnik, hep-ph/0408298. Based on talk given by J.O. Eeg at BEACH2004, 6th international conference on Hyperons, Charm and Beauty Hadrons, Illinois Institute of Technology, june 27.-july 3., Chicago, 2004.
J.O. Eeg, S. Fajfer, and A. Prapotnik, to appear.
- [15] M. Beneke, G. Buchalla, M. Neubert, C.T. Sachrajda, *Phys. Rev. Lett.* **83** (1999) 1914.
- [16] C. W. Bauer, S. Fleming, D. Pirjol, and I. W. Stewart *Phys. Rev. D* **63** (2001) 114020,
C. W. Bauer, D. Pirjol, I. W. Stewart, *Phys. Rev. D* **66** (2002) 054005,
M. Beneke and T. Feldmann, *Phys. Lett. B* **553**(2003) 267-276.
- [17] S. Weinberg, *Physica A* **96** (1979) 327.
- [18] J. Gasser and H. Leutwyler, *Ann. Phys.(N.Y.)* **158** (1984) 142, and
Nucl. Phys. B **250** (1985) 465.
- [19] See for instance: S. Bertolini, J.O. Eeg and M. Fabbrichesi, *Rev. Mod. Phys.* **72** (2000) 65-93,
and references therein.
A.J. Buras, M. Jamin *JHEP* 0401 (2004) 048, hep-ph/0306217
A. Pich, To appear in the proceedings of 32nd International Conference on High-Energy Physics (ICHEP 04), Beijing, China, 16-22 Aug 2004; hep-ph/0410215
- [20] M. Wise, Lectures given at CCAST Symp. on Particle Physics at the Fermi scale,
May 27 - June 4, 1993. Published in CCAST Symposium 1993:71-114 (QCD161:S12:1993),

- hep-ph/9306277.
- E. Jenkins, *Nucl.Phys.* **B412** (1994) 181.
- B. Grinstein, Lectures given at 6th Mexican School of Particles and Fields, Villahermosa, Mexico, 3-7 Oct 1994. Published in Mexican School 1994:122-184 (QCD161:M45:1994), hep-ph/9508227.
- C. Boyd and B. Grinstein, *Nucl. Phys.* **B442** (1995) 205.
- I. Stewart, *Nucl. Phys.* **B529** (1998) 62.
- [21] X. Ji, M. J. Musolf, *Phys. Lett.* **B 257** (1991) 409.
- D. Broadhurst, A. Grozin, *Phys. Lett.* **B 267** (1991) 105.
- [22] N. Isgur, M. Wise, *Phys. Lett.* **B 232** (1990) 527.
- [23] V. Novikov, M. Shifman, A. Vainshtein, V. Zakharov, *Fortschr. Phys.* **32** (1984) 11.
- [24] J.A.M. Vermaseren, “Symbloic Manipulation with FORM”, CAN 1991, Amsterdam. (ISBN 90-74116-01-9).
- [25] B. Grinstein and A. Falk, *Phys. Lett* **B 247** (1990) 406.
- [26] G. Amorós, M. Beneke, M. Neubert *Phys. Lett.* **B 401** (1997) 81.
- A. Czarnecki, A. Grozin, *Phys. Lett.* **B 405** (1997) 142.
- [27] A. Hiorth and J.O. Eeg, *Eur. Phys. J. direct* (2003) DOI 10.1140/epjcd/s2004-01-003-1.
- [28] See for example: G. Buchalla, A. Buras, and M. Lautenbacher, *Rev. Mod. Phys.* **68** (1996) 1125.
- [29] V. Giménez, *Nucl. Phys.* **B 401** (1993) 116.
- M. Ciuchini, E. Franco, and V. Giménez, *Phys.Lett.* **B 388** (1996) 167.
- [30] W. Kilian and T. Mannel *Phys.Lett.* **B301** (1993) 382.
- [31] D. Melikhov and N. Nikitin, *Phys. Lett.* **B 494** (2000) 229-236.
- See also: J.G. Körner, A.I. Onishchenko, A.A. Petrov, and A.A. Pivovarov, *Phys.Rev.Lett.* (2003) 91:192002 e-Print Archive: hep-ph/0306032.
- A. A. Pivovarov, hep-ph/0409257 and hep-ph/0410046.
- [32] J.M. Flynn, C.T. Sachrajda, *Adv.Ser.Direct.High Energy Phys.* **15** (1998) 402-452.
- D. Bećirević, D. Meloni, A. Retico, V. Gimenez , L. Giusti, V. Lubicz, G. Martinelli, *Nucl. Phys.* **B 618** (2001) 241-258.
- D. Bećirević, P. Boucaud, V. Gimenez, C.J.D. Lin, V. Lubicz, G. Martinelli, M. Papinutto,

- C.T. Sachrajda, Presented at 20th International Symposium on Lattice Field Theory (LATTICE 2002), Boston, Massachusetts, 24.-29. June 2002, hep-lat/0209131.
- [33] B. Grinstein, W. Kilian, T. Mannel, and M.B. Wise, *Nucl. Phys.* **B 363** (1991) 19.
R. Fleischer, *Nucl. Phys.* **B 412** (1994) 201.
- [34] N. Isgur, and M. Wise, *Phys. Rev.* **D 41** (1990) 151.
- [35] G. Ecker, J. Gasser, H. Leutwyler, A. Pich, and E. De Rafael,
Phys.Lett. **B 223** (1989) 425-432,
J. Bijnens and E. Pallante, *Mod. Phys. Lett.* **A 11** (1996) 1096-1080.
J. Bijnens, P. Gosdzinsky and P. Talavera, *Nucl. Phys.* **B 501** (1997) 495-517.
G. D'Ambrosio and J. Portoles, *Nucl. Phys.* **B 533** (1998) 494-522.
Xiao-Jun Wang and Mu-Lin Yan, hep-ph/9907321.
- [36] B. Bajc, S. Fajfer, and R.J. Oakes, *Phys. Rev.* **D 53**(1996) 4957.
- [37] J. Charles, A. Le Yaouanc, L. Oliver, O. Pene, and J.C. Raynal, *Phys.Rev.* **D60** (1999) 014001,
M. Beneke and T. Feldmann, *Nucl. Phys.* **B 592** (2001) 3-34.
- [38] U. Agleatti, G: Corbò, and L. Trentadue , *Int. J. Mod. Phys.* **A 14** (1999) 1769.
U. Agleatti and G: Corbò, *Int. J. Mod. Phys.* **A 15** (2000) 363.